

EXHIBIT 9

Applicants: Eran Blaugrund et al.

Serial No.: 10/712,958

Filed: November 13, 2003

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 538 134 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
12.03.1997 Bulletin 1997/11

(51) Int Cl.⁶: **C07C 211/42, A61K 31/135**

(21) Application number: 92402821.0

(22) Date of filing: 15.10.1992

(54) **Mono-fluorinated derivatives of N-propargyl-1-aminoindan and their use as inhibitors of monoamine oxidase**

Monofluorierte Derivate von N-Propargyl-1-Aminoindan und deren Verwendung als Monoamino-Oxidaseinhibitoren

Dérivés mono-fluorés de N-propargyl-1-aminoindane et leur usage comme inhibiteurs de monoamine oxydase

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
PT SE

(30) Priority: 16.10.1991 IL 99759

(43) Date of publication of application:
21.04.1993 Bulletin 1993/16

(73) Proprietor: **TEVA PHARMACEUTICAL
INDUSTRIES, LTD.**
Jerusalem 91010 (IL)

(72) Inventors:
• Sterling, Jeff
Jerusalem (IL)
• Levy, Ruth
Tel-Aviv 69415 (IL)

- Veinberg, Alex
Rehovot 76281 (IL)
- Goldenberg, Willy
Jerusalem 93461 (IL)
- Finberg, John
Tivon 36000 (IL)
- Youdim, Musa
Haifa 32763 (IL)
- Gutman, Arleh
Ahuza, Haifa 34752 (IL)

(74) Representative: **Le Roux, Martine et al**
Cabinet Beau de Loménie
158, rue de l'Université
75340 Paris Cédex 07 (FR)

(56) References cited:
EP-A- 0 436 492 WO-A-85/05617
GB-A- 1 003 686

Applicants: Eran Blaugrund et al.
Serial No.: 10/712,958
Filed: November 13, 2003
Exhibit 9

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

The present invention is in the field of selective irreversible inhibitors of the enzyme monoamine oxidase (hereinafter MAO) and relates to novel propargylamine compounds which are selective irreversible inhibitors of the B-form of the monoamine oxidase enzyme (hereinafter, MAO-B). The invention also relates to pharmaceutical compositions containing these propargylamine compounds which are particularly useful for the treatment of Parkinson's disease, memory disorders and dementia of the Alzheimer type (DAT), depression, and hyperactive syndrome in children.

Parkinson's disease is widely considered to be the result of degradation of the pre-synaptic dopaminergic neurons in the brain, with a subsequent decrease in the amount of the neurotransmitter dopamine, that is being released. Inadequate dopamine release, therefore, leads to the onset of voluntary muscle control disturbances symptomatic of Parkinson's disease.

Various procedures for treating Parkinson's disease have been established and are currently in widespread use, for example, the administration of L-Dopa, which is a precursor of dopamine, together with a decarboxylase inhibitor, such as L-carbidopa or benzerazide. The decarboxylase inhibitor protects the L-Dopa molecule from peripheral decarboxylation and thus ensures L-Dopa uptake by the remaining dopaminergic neurons in the striatum of the brain. Here the L-Dopa is converted into dopamine resulting in increased levels of dopamine in these neurons. In response to physiological impulses these neurons are therefore capable of releasing larger amounts of dopamine, the quantity of which approximates the normal required levels. This treatment therefore alleviates the symptoms of the disease and contributes to the well-being of the patients.

However, this L-Dopa treatment has its drawbacks, the main one being that its effectiveness is optimal only in the first few years following the onset of treatment. After this initial period the clinical response is diminished and is accompanied by adverse side effects which include dyskinesia, fluctuation in efficacy throughout the day ("on-off effect") and psychiatric symptoms such as confusional states, paranoia and hallucinations. This fall-off in the effect of L-Dopa treatment is attributed to a number of factors, including the natural progression of the disease, alteration in dopamine receptors as a consequence of increased dopamine production or increased levels of dopamine metabolites, and pharmacokinetic problems of L-Dopa absorption (reviewed by Youdim et al., Progress in Medicinal Chemistry, Vol. 21, Chapter 4, pp. 138-167 (1984), Eds. Ellis and West, Elsevier, Amsterdam).

In order to overcome the drawbacks of the L-Dopa treatment, various treatments have been devised in which L-Dopa is combined with MAO inhibitors, with the aim of reducing the metabolic breakdown of the newly formed dopamine (see for example, U.S. 4,826,875).

MAO exists in two forms known as MAO-A and MAO-B which have selectivity for different substrates and inhibitors. For example, MAO-B metabolises more efficiently substrates such as 2-phenylethylamine and is selectively and irreversibly inhibited by (-)-deprenyl (as described below).

It should be noted, however, that combining L-Dopa with an inhibitor of both MAO-A and MAO-B is undesirable leading to adverse side effects related to an increased level of catecholamines throughout the neuraxis. Furthermore, complete inhibition of MAO is also undesirable as it potentiates the action of sympathomimetic amines such as tyramine leading to the so-called "cheese effect" (reviewed by Youdim et al., Handbook of Experimental Pharmacology, Vol. 90, Chap. 3 (1988) Eds. Trendelenburg and Weiner, Springer-Verlag). As MAO-B was shown to be the predominant form of MAO in the brain, selective inhibitors for this form were thus considered to be a possible way for achieving a decrease in dopamine breakdown on the one hand, together with a minimization of the systemic effects of total MAO inhibition, on the other.

One of these selective MAO-B inhibitors, (-)-deprenyl, has been extensively studied and has been used as an MAO-B inhibitor to augment L-Dopa treatment. This treatment with (-)-deprenyl is generally favourable, not causing the "cheese effect" at doses causing nearly complete inhibition of MAO-B (Elsworth et al., Psychopharmacology, 57, 33 (1978). Furthermore, addition of (-)-deprenyl to a combination of L-Dopa and decarboxylase inhibitor to Parkinson's patients leads to improvements in akinesia and overall functional capacity as well as the elimination of "on-off" type fluctuations (reviewed by Birkmayer & Riederer in "Parkinson's Disease" pp. 138-149, Springer-Verlag (1983)).

Thus, (-)-deprenyl enhances and prolongs the effect of L-Dopa and permits a lowering of the dosage of L-Dopa whereby the adverse effects of L-Dopa treatment are limited.

A more potent selective inhibitor of MAO-B than the racemic mixture is the optical isomer R-(+)-N-propargyl-1-aminoindan. HCl [R-(+)-PAI.HCl] which is more selective in vivo and in vitro as described in EP-A-0 436 492.

However it is highly desirable to further increase the selectivity of MAO inhibitor, inhibiting preferably MAO-B and not MAO-A, thus minimizing the side effects caused by the inhibition of MAO-A.

The compounds of the present invention were found to have a surprisingly high degree of selectivity in vitro, inhibiting preferably MAO-B over MAO-A.

The compounds of the present invention are mono-fluoro derivatives of N-propargyl-1-aminoindan stereoisomers and salts thereof.

US 3,513,244 claims generically and specifically a large number of secondary and tertiary aminoindans which are


stated to have hypotensive activity. There is no disclosure of the specific compounds of the present invention, and there is no evidence that the mono-fluorinated derivatives of 1-propargylaminoindan of the present invention have ever been synthesised and characterised. Furthermore, the secondary and tertiary aminoindans of US 3,513,244 are stated to have hypotensive activity and there is no teaching that any components of the group have any as MAO-B inhibitor activity.

GB 1,003,686 discloses a group of benzocycloalkane compounds in which the cycloalkane has from five to seven ring members and is substituted by an N-(alkynylalkyl)amino group. This patent mentioned the possibility of substituting the aromatic portion of the benzocycloalkane ring system by one or more halogen atoms. Although the compounds of the present invention are generically included in the group, they are not specifically disclosed. The compounds claimed in GB 1,003,683 are stated to be monoamine oxidase inhibitors in general, but there is no disclosure of any activity by which MAO-B is inhibited, selectively in preference over MAO-A.

The mono-fluorinated N-propargyl-1-aminoindans of the present invention are surprisingly more potent selective MAO-B inhibitors than any of the other species of the groups disclosed in both US 3,513,244 and GB 1,003,686, enabling the inhibition of MAO-B at lower concentrations. This high potency and selectivity in vivo is a unique and surprising property of the mono-fluorinated N-propargyl-1-aminoindan compounds and their salts according to the invention, not shared by other halogenated N-propargyl-1-aminoindans.

The MAO-B inhibitors of the present invention can also be used for the treatment of patients with dementia of the Alzheimer type (DAT), and the treatment of patients with depression and for treatment of hyperactive syndrome in children. Other MAO-B inhibitors have been used in the past for the treatment of patients suffering from these diseases (Tariot et al., *Psychopharmacology*, 91, 489-495, 1987), J. Mendlewich and M.B.H. Youdim (*Brit. J. Psychiat.*, 142, 508-511, 1983, Perenyi A., et al., PCT/HU89/00044, 17 August 1988).

The present invention provides novel mono-fluorinated derivatives of 1-propargyl aminoindans of the formula



and pharmaceutically acceptable addition salts thereof.

The compounds of formula (I) may be racemic mixtures or optically pure enantiomers. Specific compounds of use in the present invention include

4-fluoro-N-propargyl-1-aminoindan
5-fluoro-N-propargyl-1-aminoindan
6-fluoro-N-propargyl-1-aminoindan
and optically pure enantiomers thereof.

A preferred compound of the invention is 6-fluoro-N-propargyl-1-aminoindan and particularly preferred compound is the enantiomer (+)-6-fluoro-N-propargyl-1-aminoindan.


The present invention also relates to pharmaceutical compositions comprising compounds of formula (I) together with pharmaceutically acceptable carriers and/or excipients and/or diluents. The pharmaceutical compositions may be adapted for oral, rectal, parenteral, topical or transdermal administration. Suitable forms for oral administration include tablets, compressed or coated pills, dragees, sachets, hard or soft gelating capsules, syrups and suspensions. Suitable forms for parenteral administration include ampoules or vials which additionally contain an aqueous or nonaqueous solution or emulsion. Compositions adapted for rectal administration include suppositories with hydrophilic and/or hydrophobic vehicles.

The pharmaceutical compositions may be in dosage unit forms preferably containing 1-20 mg of the compound according to formula (I). The pharmaceutical compositions may additionally comprise levodopa and a decarboxylase inhibitor, such as L-carbidopa or benserazide. Preferably the pharmaceutical composition will comprise 1-20 mg of the compound according to formula (I), 50-250 mg levodopa and 10-25 mg of L-carbidopa. Another preferable pharmaceutical composition will comprise 2-10 mg of the compound according to formula (I), 50-250 mg levodopa and 12.5-50 mg benserazide.

The present invention further relates to the use of the compound of formula (I) for manufacturing a pharmaceutical composition for the treatment of human patients suffering from Parkinson's disease, memory disorders, dementia of

the Alzheimer type and hyperactive syndrome in children.

The invention also relates to processes for the preparation of the compounds of formula (I). The racemic mixture of the compound of formula (I) can be prepared by reacting aryl fluorinated 1-chloro or 1-bromoindans, with propargylamine. Alternatively, these racemates may be prepared by reacting propargylamine with substituted 1-indanones to form the corresponding imines, followed by reduction of the carbon-nitrogen double bond of the imine with a suitable agent, such as sodium borohydride. Another method of preparation of the racemic mixture of the compound of formula (I) comprises the reaction of a racemic fluorinated 1-aminoindan of the formula:




with propargyl bromide or propargyl chloride in the presence of an organic or inorganic base, optionally in the presence of a suitable solvent and, if desired, converting the resulting free base of formula I into a pharmaceutically acceptable acid addition salt thereof.

The racemic mixture of (+) and (-) enantiomers of fluorinated 1-aminoindan may be prepared by chemical reduction of corresponding fluorinated oximes, e.g., with Zn in acetic acid or by catalytic hydrogenation. Fluorinated indan-1-one may be prepared, e.g., by Friedel-Crafts cyclization of fluorinated dihydrocinnamic chloride using aluminium chloride or other Lewis acids as condensing agents. Fluorinated dihydrocinnamic chlorides may be prepared as known per se.

The enantiomers of the compounds of formula (I) may be obtained by optical resolution of racemic mixtures of (+) and (-) enantiomers of compounds of formula (I). Such a resolution can be accomplished by conventional resolution methods, well known to a person skilled in the art, such as those described in "Enantiomers, Racemates and Resolutions" by J. Jacques, A. Collet and S. Willen, Pub. John Wiley & Sons, NY 1981. For example, the resolution may be carried out by preparative chromatography on a chiral column. Another suitable resolution method is the formation of diastereomeric salts with a chiral acid such as tartaric, maleic, mandelic acid or N-acetyl derivatives of amino acids, such as N-acetyl leucine, followed by recrystallization to isolate the diastereomeric salt of the desired enantiomer.

In accordance with this invention, the (+) enantiomers of the compounds formula (I) can be prepared directly from the optically active (-) enantiomers of fluorinated 1-aminoindans of the formula



by reaction with propargyl bromide or propargyl chloride in the presence of an organic or inorganic base and optionally in the presence of a suitable solvent and if desired converting the resulting (+) enantiomeric free base of formula I into a pharmaceutically acceptable acid salt thereof.

Suitable organic or inorganic bases for use in the above reaction are, e.g., triethylamine, pyridine, alkali metal carbonates or bicarbonates etc. If the reaction is conducted in the presence of a solvent, this may be chosen from, e.g., toluene, methylene chloride and acetonitrile. A preferred method of preparation of the aforementioned compounds is the reaction between fluorinated (-)-1-aminoindans with propargyl chloride using potassium carbonate as a base and acetonitrile as solvent.


The above described reaction between fluorinated 1-aminoindans generally results in a mixture of unreacted primary amines, the desired secondary amines and tertiary amines, namely the N,N-bispropargylamino products. The desired secondary amines, i.e. fluorinated N-propargyl-1-aminoindans, can be separated by conventional separation methods including but not limited to chromatography, distillation and selective extraction.

The compounds of formula (II) herein as well as their acid addition salts are novel. These compounds, which are

intermediates in the preparation of compounds of formula (I) herein, are provided as racemates and as enantiomers, preferably as (-) enantiomers.

Alternatively, the fluorinated (-)-1-aminoindans may be prepared by reacting fluorinated 1-indanone with an optically active amine, followed by reduction of the carbon-nitrogen double bond of the resulting imine by hydrogenation over a suitable catalyst such as platinum oxide, Raney nickel, or by chemical reduction, for example, with sodium borohydride. Suitable optically active amines are, for example, one of the antipodes of alpha-phenylethylamine or an ester of an amino acid, such as phenylalanine. The benzylic N-C bond may be cleaved by hydrogenolysis.

Fluorinated (-) aminoindans may also be prepared by enzyme mediated selective acylation of a racemic mixture of a compound of the formula:



preferably using the enzyme subtilisin A isolated from *Bacillus licheniformis* in 3-methyl-3-pentanol with trifluoroethyl butyrate as the acylating agent thereby forming a mixture comprising an acylated (+) enantiomer of fluorinated-1-aminoindan and a non-acylated (-) enantiomer of fluorinated aminoindan then the resulting fluorinated (-)-1-aminoindan may be readily separated from the corresponding (+)-amide by chromatography, distillation, selective extraction, or conversion of the free base into a suitable acid addition salt and its fractional recrystallisation.

Finally the (+)-fluoro-N-propargyl-1-aminoindan is prepared by reacting the separated (-) fluoro-1-aminoindan with propargyl bromide or propargyl chloride in the presence of an organic or inorganic base, optionally in the presence of a suitable solvent, and isolating the (+) fluoro-N-propargyl-1-aminoindan formed as the free base or a pharmaceutically acceptable acid addition salt thereof.

Additional methods for preparing fluorinated (-)-1-aminoindans are the reduction, as described above, of indan-1-one oxime ethers, wherein the alkyl portions of the ether contains an optically pure chiral centre. Alternatively, a non-chiral derivative of fluorinated indan-1-ones containing a carbon-nitrogen double bond, such as an imine or oxime, can be reduced with a chiral reducing agent, e.g., a complex of lithium aluminium hydride and ephedrine.

Pharmaceutically acceptable acid addition salts of the compounds of the present invention may be prepared by reacting according to conventional methods the free base forms of the compounds of formula (I) with the desired acids in the presence of a suitable solvent. Similarly, an acid addition salt may be converted to the free base form or directly to another acid addition salt by methods well known to those skilled in the art.

Thus, the present invention also provides a method for the preparation of a (-) enantiomer of the formula (III) comprising:

- a) selectively acylating a racemic mixture of a compound of the formula (II), with a suitable acylating agent in the presence of the enzyme subtilisin A isolated from *Bacillus licheniformis* thereby forming a mixture comprising an acylated (+) enantiomer of fluorinated-1-aminoindan and a non-acylated (-) enantiomer of fluorinated-1-aminoindan,
- b) separating the (-) halogenated-1-aminoindan from the acylated (+) fluorinated-1-aminoindan by chromatography, distillation, selective extraction or conversion of the free base into a suitable acid addition salt and its recrystallisation.

The present invention will not be specifically described in the following Examples to which it is not limited.

Example 1: 5-Fluoro-1-aminoindan

A solution of 3-fluorobenzaldehyde (10g), malonic acid (15.6 g) and piperidine (0.7 ml) in pyridine (35 ml) was heated at reflux for 3.5 hours. After cooling to room temperature, the reaction mixture was added to a mixture of ice water (210 ml) and cone. hydrochloric acid (56 ml). The precipitate was isolated by filtration and recrystallised from methanol/water to give 3-fluorocinnamic acid (11.3 g).

A mixture of 3-fluorocinnamic acid (11 g), sodium hydroxide (2.7 g), 3% sodium amalgam (220 g) in 340 ml of water was heated overnight at 70-80°C. The aqueous solution was decanted from the mercury residue and acidified

with conc. hydrochloric acid (35 ml) to pH 2, cooled at 0°C and the resulting precipitated 3-fluorohydrocinnamic acid (10 g) isolated by filtration.

A solution of 3-fluorohydrocinnamic acid (28 g) in thionyl chloride (90 ml) was heated at reflux for 3 hours. 3-Fluorohydrocinnamoyl chloride (27 g) was isolated by distillation (150-110°C @ ca 20 mm.Hg).

A solution of 3-fluorodihydrocinnamoyl chloride (15.4 g) in carbon disulfide (78 ml) was added dropwise to a cold (-5°C) mixture of aluminium chloride (13.9 g) in carbon disulfide (300 ml) for 30 minutes. During this time and for 30 minutes the mixture was allowed to warm to room temperature (1 hr), then it was heated to reflux (1 hr) and finally the volatiles were removed by distillation under reduced pressure. The residue was dissolved in methylene chloride (400 ml) and washed successively with 10% aqueous sodium hydroxide and water. After drying (magnesium sulfate) and solvent removal in vacuo, 5-fluoroindanone (2 g) was isolated by recrystallisation from hexane.

A solution of 5-fluoroindanone (10 g), hydroxylamine hydrochloride (9 g), potassium carbonate (23 g) and water (3.5 ml) in 96% ethanol (40 ml) was heated to reflux, cooled to ca 45°C and additional hydroxylamine hydrochloride (4.5 g) and potassium carbonate (11.5 g) were added and refluxed for an additional 30 min. At this time the mixture was poured into cold water (300 ml) and stirred in an ice bath for one hour; 5-fluoroindanone oxime (9.8 g) was isolated by filtration.

Zinc powder (20.2 g) was added during three hours to the stirred solution of the above oxime (20.2 g) in acetic acid (200 ml) while maintaining the temperature at 25-35°C. The resulting mixture was stirred an additional 12 hours at room temperature. The solids were removed by filtration and the filtrate concentrated in vacuo. The concentrated filtrate was partitioned between water and toluene and the aqueous phase adjusted to pH 12 with ammonium hydroxide and extracted with toluene. Solvent removal in vacuo gave 5-fluoro-1-aminoindan (17.8 g) as a colored oil. IR: 690, 748, 815, 864, 929, 1126, 1244, 1315, 1377, 1433, 1454, 1485, 1595, 1614, 2800, 3000, 3300 cm⁻¹. NMR: 1.81, 2.4, 2.8, 6.8, 7.2; m/e 151.0786 C₉H₁₀NF;

Example 2: 4-fluoro-1-aminoindan

The title compound was prepared in 47% overall yield according to the procedure of Example 1 except that 2-fluorobenzaldehyde was used instead of 3-fluorobenzaldehyde.

IR: 708, 780, 1170, 1242, 1290, 1377, 1472, 1587, 1624, 2800, 2950, 3200 cm⁻¹;

NMR: 1.7, 2.1, 2.7, 2.9, 3.1, 4.3, 6.8, 7.0, 7.1;

m/e: 151.0782 (C₉H₁₀NF)

Example 3: 6-fluoro-1-aminoindan

The title compound was prepared in 29% overall yield according to the procedure of Example 1 except that 4-fluorobenzaldehyde was used instead of 3-fluorobenzaldehyde.

IR: 694, 740, 770, 812, 844, 870, 912, 1130, 1163, 1253, 1441, 1483, 1597, 1612, 2800, 3000, 3300 cm⁻¹;

NMR: 1.5, 2.1, 2.8, 6.8, 7.1; m/e 151.0774 C₉H₁₀NF

Example 4: (-)-6-Fluoro-1-aminoindan

A solution of 6-fluoro-1-aminoindan (4.2 g) in methanol (10 ml) was added to a heated clear solution of L-N-acetyl-methyl-3,4-dimethoxyphenylalanine (7.8 g) in methanol (30 ml). The mixture was stirred with cooling. The product was crystallised from methanol with carbon black (0.5 g) two times and checked by HPLC (chiral column).

The title compound obtained by extraction from aqueous 10% sodium hydroxide with methylene chloride, drying and evaporation. [α]_D-7.9° (c 2%, EtOH).

The spectral properties were identical to the compound of Example 3.

Example 5: (+)-6-Fluoro-1-aminoindan

The title compound was crystallised from mother liquors after separation of (-)-antipode by the procedure of Example 4, and repeated crystallisation from the same solvent. The title compound was obtained by extraction from aqueous 10% sodium hydroxide with methylene chloride, drying and evaporation. The spectral properties were identical to the compound of Example 3.

Example 6: (-)-4-Fluoro-1-aminoindan

The title compound was prepared in 50% yield according to the procedure of Example 4 except that 4-fluoro-1-aminoindan was used instead of 6-fluoro-1-aminoindan. The spectral properties were identical to the compound of

Example 2.

Example 7: (-)-5-Fluoro-1-aminoindan

The title compound was prepared in 25% yield according to the procedure of Example 4 except that 5-fluoro-1-aminoindan was used instead of 6-fluoro-1-aminoindan. The spectral properties were identical to the compound of Example 1.

Example 8: (-)-6-Fluoro-1-aminoindan

A solution of 20 grams of racemic 6-fluoro-1-aminoindan and 60 ml of trifluoroethyl butyrate in 400 ml of 3-methyl-3-pentanol was treated with 2 g of subtilisin A. The resulting suspension was shaken on an orbital shaker at 200 rpm at 40°C for 72 hours at which time the enzyme was removed by filtration, and the filtrate was extracted with 1M aqueous hydrochloric acid. The aqueous layer was back extracted with methylene chloride and freeze dried to yield 10 grams of the title compound as the HCl salt. The free base was obtained by suspending the HCl salt in aqueous 10% sodium hydroxide, extracting with methylene chloride, drying and evaporation. The free base so obtained was of 99% optical purity with spectroscopic properties identical to those of the compound from Example 4.

Example 9: (-)-5-Fluoro-1-aminoindan

The title compound was obtained in 99% optical purity by the method of Example 8 using racemic 5-fluoro-1-aminoindan instead of 6-fluoro-1-aminoindan.

Example 10: (-)-4-Fluoro-1-aminoindan

The title compound was obtained in 99% optical purity by the method of Example 8 using racemic 4-fluoro-1-aminoindan instead of 6-fluoro-1-aminoindan.

Example 11: 5-Fluoro-N-propargyl-1-aminoindan hydrochloride

A mixture of 5-fluoro-1-aminoindan (6.2 g), potassium carbonate (7.4 g) and acetonitrile (54 ml) were heated under nitrogen to 60°C. Propargyl chloride (3.3 g) was added and heating was continued overnight. Acetonitrile was evaporated and products were distributed between methylene chloride and 10% sodium hydroxide in water.

The free base of the title compound was isolated by flash column chromatography on silica gel.

An ethereal solution of this base was treated with hydrogen chloride gas and the title compound (2.8 g) was isolated by filtration.

M.p. 178.4°C (decomp.); IR (KBr) 899, 1216, 1437, 1491, 2359, 2928, 3287 cm^{-1} . Anal. calcd. for $\text{C}_{12}\text{H}_{15}\text{ClFN}$: % C 63.86, H 5.81, N 6.21, Cl 15.71; found % C 63.38, H 5.69, N 6.33, Cl 15.00.

Example 12: 4-Fluoro-N-propargyl-1-aminoindan hydrochloride

The title compound was prepared in 29% yield according to the procedure of Example 11 except that 4-fluoro-1-aminoindan was used instead of 5-fluoro-1-aminoindan.

M.p. 191.3°C (decomp.); IR (KBr) 771, 1045, 1248, 1473, 1586, 2400, 2653, 2719, 2949, 3200 cm^{-1} ; m/e 190. Anal. calcd. for $\text{C}_{12}\text{H}_{15}\text{ClFN}$: % C 63.86, H 5.81, N 6.21, Cl 15.71; found % C 63.66, H 5.58, N 6.15, Cl 15.54.

Example 13: 6-Fluoro-1-propargylaminoindan

The title compound was prepared in 44% yield according to the procedure of Example 11 except that 6-fluoro-1-aminoindan was used instead of 5-fluoro-1-aminoindan. The spectral properties were identical to the compound of Example 16.

Example 14: (+)-4-Fluoro-N-propargyl-1-aminoindan hydrochloride

The title compound was prepared in 70% yield according to the procedure of Example 11 except that (-)-4-fluoro-1-aminoindan was used instead of 5-fluoro-1-aminoindan.

M.p. 218.9°C (decomp.); $[\alpha]_D^{25} + 10.7$ (c, 0.4%, H_2O); NMR (D_2O) 2.35 m, 2.65 m, 3.00 m, 3.93 s, 4.8 s, 5.00 m, 7.15 m, 7.33 m ppm; m/e 189.09;

IR (KBr) 771, 1248, 1583, 2438, 2655, 2718, 2849, 2945, 3239 cm^{-1} ; Anal. calcd. for $\text{C}_{12}\text{H}_{15}\text{ClFN}$: % C 63.86, H 5.81, N 6.21, Cl 15.71; found % C 64.01, H 5.74, N 6.05, Cl 15.80.

Example 15: (+)-5-Fluoro-N-propargyl-1-aminoindan hydrochloride

The title compound was prepared in 50% yield according to the procedure of Example 11 except that (-)-5-fluoro-1-aminoindan was used instead of 5-fluoro-1-aminoindan.

NMR (CDCl_3) 2.35 m, 2.8 m, 3.3 m, 3.6 s, 4.6 d, 6.8 m, 7.6 m ppm; m/e 189.09; IR (KBr) 690, 713, 833, 945, 1016, 1238, 1252, 1420, 1444, 1487, 1600, 1625, 2100, 2400, 2500, 2900, 3250 cm^{-1} ; $[\alpha]_D^{25} +23.5^\circ$ (c, 0.2%, H_2O); M.p. 192°C (decomp.);

Anal. calcd. for $\text{C}_{12}\text{H}_{15}\text{ClFN}$: % C 63.86, H 5.81, N 6.21, Cl 15.71; found % C 64.52, H 5.68, N 6.35, Cl 15.47.

Example 16: (+)-6-Fluoro-N-propargyl-1-aminoindan hydrochloride

The title compound was prepared in 46% yield according to the procedure of Example 11 except that (-)-6-fluoro-1-aminoindan was used instead of 5-fluoro-1-aminoindan.

M.p. 220.8°C (decomp.); $[\alpha]_D^{25} +18.1^\circ$ (c, 0.3%, H_2O);

IR (KBr) 694, 735, 820, 880, 1042, 1131, 1173, 1228, 1246, 1256, 1365, 1460, 1492, 1584, 1599, 2128, 2438, 2946, 3064, 3224 cm^{-1} . Anal. calcd. for $\text{C}_{12}\text{H}_{15}\text{ClFN}$: % C 63.86, H 5.81, N 6.21, Cl 15.71;

found % C 64.39, H 5.54, N 6.21, Cl 15.71.

Example 17: (-)-6-Fluoro-N-propargyl-1-aminoindan hydrochloride

The title compound was prepared in 61% yield according to the procedure of Example 11, except that (+)-6-fluoro-1-aminoindan was used instead of 5-fluoro-1-aminoindan.

M.p. 220°C; $[\alpha]_D^{25} -17.7^\circ$; IR(KBr) 828, 1130, 1228, 1493, 1597, 2440, 2633, 2945, 3229 cm^{-1} .

Example 18: 6-Chloro-N-propargyl-1-aminoindan hydrochloride

The title compound was prepared in 33.1% yield according to the procedure of Example 11 except that 6-chloro-1-aminoindan was used instead of 5-fluoro-1-aminoindan.

M.p. 191.7°C (decomp.); NMR (D_2O) 2.3 m, 2.5 m, 3.1 m, 3.93 s, 4.9 d, 5 m ppm; m/e 205.0658 and 207.0640.

IR (KBr) 760, 812, 1091, 1259, 1472, 1578, 1599, 1651, 1700, 2900, 3300 cm^{-1} ; Anal. calcd. for $\text{C}_{12}\text{H}_{13}\text{NCl}_2$: % C 59.52, H 5.41, N 5.78, Cl 29.28; found % C 59.80, H 5.33, N 5.61, Cl 29.11.

Example 19: 4,6-Difluoro-N-propargyl-1-aminoindan hydrochloride

The title compound was prepared in 61.8% yield according to the procedure of Example 11 except that 4,6-difluoro-1-aminoindan was used instead of 5-fluoro-1-aminoindan.

M.P. 218.8°C (decomp.); NMR (D_2O) 1.16, 2.36 m, 2.63 m, 3.02 m, 3.98 s, 7.01 s, 7.18 d; m/e 207.0817;

IR (KBr) 851, 872, 980, 995, 1055, 1113, 1335, 1447, 1462, 1487, 1584, 1601, 1630, 2400, 2700, 2950, 320 cm^{-1} ; Anal. calcd. for $\text{C}_{12}\text{H}_{12}\text{NClF}_2$: % C 59.15, H 4.96, N 5.75, Cl 14.55; found % C 59.41, H 5.19, N 5.61, Cl 13.98.

Example 20: Pharmaceutical formulation

A typical pharmaceutical preparation can be prepared as follows:
Each tablet contains:

6-fluoro-N-propargylaminoindan hydrochloride	3.0 mg
Pregelatinized starch	11.0 mg
Starch	48.5 mg
Microcrystalline cellulose	80.0 mg
Ethylcellulose	1.0 mg
Talc	1.5 mg
Tablet weight:	145.0 mg

Blend the excipients and the active granulate with isopropyl alcohol. Dry the mixture, blend with the talc and com-

press into tablets.

Similar formulations can be prepared by one skilled in the art for each of the compounds of the invention.

Example 21: Tablet Composition

Each tablet contains:

6-fluoro-N-propargyl-1-aminoindan-hydrochloride	5.0 mg
Levodopa	100.0 mg
Carbidopa	25.0 mg
Pregelatinised starch	24.0 mg
Starch	40.0 mg
Microcrystalline cellulose	49.5 mg

Alcohol USP added as required to granulation.

Similar formulations can be prepared by one skilled in the art for each of the compounds of the invention.


Example 22: Inhibition of MAO Activity In-Vitro

Experimental Protocol:

These experiments were performed essentially according to the protocol of K.F. Tipton and M.B.H. Youdim in CIBA Symposium No. 39, Editors: G.E.W. Wolstenholme and J. Knight, Elsevier, Amsterdam, 1976, pp. 393-403. The MAO enzyme source was a homogenate of rat brain in 0.3 M sucrose which was centrifuged at 600 g for 15 minutes. The supernatant was diluted appropriately in 0.05M phosphate buffer and preincubated with serial dilutions of compounds of interest which are listed below for 20 minutes at 37°C. ¹⁴C-labeled substrates (2-phenylethylamine, hereinafter PEA; 5-hydroxytryptamine, hereinafter 5-HT) were then added and the incubation continued for a further 20 minutes (PEA) or 30-45 minutes (5-HT). Substrate concentrations used were 20 µM (PEA) or 1mM (5-HT). In the case of PEA the enzyme concentration was chosen so that not more than 10% of the substrate was metabolised during the course of the reaction. The reaction was then stopped by the addition of tranylcypromine (to final concentration of 1 mM) and the incubate filtered over a small column of Amberlite CG-50; buffered to pH 6.3. The column was washed with 1.5 ml water, the eluates pooled and the radioactive content determined by liquid scintillation spectrometry. Since the amine substrates are totally retained on the column, radioactivity in the eluate indicates the production of neutral and acidic metabolites formed as a result of MAO activity. Activity of MAO in the sample was expressed as a percentage of control activity in the absence of inhibitors after subtraction of appropriate blank values. The activity determined using PEA as a substrate is referred to as MAO B activity, and that determined using 5-HT as a substrate is referred to as MAO A activity. The inhibitory activity of each of the compounds was examined separately in vitro and is shown in Table 1. The results shown in Table 1 demonstrate that the fluorinated propargylaminoindans of the invention are surprisingly more selective MAO-B inhibitors than the non-halogenated parent compound. The preferred compound of the invention, (+)-6-F-propargyl-N-aminoindan is much more selective than the parent non-fluorinated compound and more selective than the other fluorinated derivatives.

TABLE 1

IC₅₀ Values (μM) for Inhibition of MAO-A and MAO-B
in Brain Homogenates



Compound Substituent R	MAO-B	MAO-A	Selectivity MAO-A/MAO-B
4-F	0.0058	0.5	86
(+)-4-F	0.0052	0.34	65
5-F	0.006	0.7	116
(+)-5-F	0.0057	0.14	24
6-F	0.0062	4.00	645
(+)-6-F	0.00022	2.8	1244
(-)-6-F	12	40	3
6-Cl	0.0077	6.9	896
(+)-4,6-Difluoro	0.5	0.8	2
H	0.003	0.073	24

IC₅₀ the concentration that caused 50% inhibition in MAO activity

Example 23: Inhibition of MAO Activity Ex-vivo: Acute Treatment


Experimental Protocol:

Rats (male Sprague-Dawley derived) weighing 250±20 g were treated with the desired compound by intraperitoneal injection (ip) or oral gavage (po) and decapitated 2 hours later. Groups of three rats were used for each dose level of the compound and MAO activity was determined in the brain and liver using the general technique described above. The amount of protein in each incubation was determined using the Folin-Lowry method, and enzyme activity was calculated as nmol substrate metabolised per hour incubation for each mg protein. Activity of MAO in tissues from animals treated with inhibitors was expressed as a percentage of the enzyme activity in a group of control animals which were administered the vehicle alone (water in the case of oral gavage, or 0.9% saline in the case of intraperitoneal injection) and killed as above. The results are presented in Table 2.

The ex-vivo experiment demonstrates that the monofluorinated derivatives of 1-propargylaminoindan of the present invention are more potent and selective in the inhibition of MAO-B as compared to MAO-A than the 6-chloro derivatives.

TABLE 2

Ex-vivo MAO Inhibitory Activity



ED-50 BRAIN (mg/kg)

ED-50 LIVER (mg/kg)

Compound Substituent R	MAO -B	MAO- A	MAO-A/ MAO-B	MAO -B	MAO- A	MAO-A/ MAO-B
4-F	0.22	1.4	6	0.13	2.4	18.5
(+)-4-F	0.33	>1.0	>3	0.17	>1.0	>5.9
5-F	0.4	2.3	6	0.06	2.2	36.7
(+)-5-F	0.07	>0.1	>1.4	0.07	>0.1	>1.4
6-F	0.13	22	170	ND	ND	ND
(+)-6-F	0.14	21.3	152	0.13	5.2	40
(-)-6-F	0.45	23	51	<0.5	ND	ND
6-Cl	1.7	10	5.8	ND	ND	ND
(+)-4,6- Difluoro	3.7	2.9	1	ND	ND	ND
H	0.07	1.2	17	0.06	5	83

ND - Not determined

ED-50 - The effective dose (mg/kg) that caused 50 %
inhibition in MAO activity

MAO -A/MAO-B -This is a measure of selectivity

Example 24: Tyramine Potentiation in Vivo

Experimental protocol:

Male rats weighing 310 ± 20 gr (5-10 per group) were treated with the desired compound by oral administration (gavage) of 5mg/kg.

On the previous day a tail artery cannulation was performed to enable direct blood pressure measurements. On the experimental day the cannulae were connected to pressure transducers. The animals were unrestrained and were

kept in a glass sided open top tank during the pressure measurements. Tyramine was administered orally by gavage in increasing doses (5, 10 and 20mg/kg), allowing 30 minutes between doses. The desired compound was administered 30 minutes after the last dose of tyramine. Tyramine was administered again, in increasing doses, as described above, one hour after the desired compound was administered. Tyramine alone caused increase of blood pressure which was potentiated by the compounds.

The tyramine potentiating effect was calculated as a ratio between the areas under the curve of blood pressure measured with tyramine alone (20mg/kg) and that measured after administration of the desired compounds and tyramine.

Results:

The results are presented in Table 3. The three fluorinated propargyl aminoindans tested caused certain potentiation of tyramine effect on blood pressure at this high dose of 5mg/kg. (+)-6-F-N-propargyl-1-aminoindan has the lowest potentiating effect as compared to the (+)-5-F and (+)-4-F derivatives.

The results of this experiment indicate that the MAO-B selectivity of the fluorinated derivatives and especially of their (+) enantiomers prevent the tyramine potentiating effect ("cheese effect") caused by the non selective MAO-inhibitors. This tyramine potentiating effects is one of the major reasons which prevented the clinical use of non-selective MAO-inhibitors. The preferred compound in this regard is (+)-6-F-N-propargyl-1-aminoindan which does not cause significant potentiation of tyramine at a high dose of 5mg/kg body weight. The ED₅₀ for brain MAO-B inhibition in rat is 0.14mg/kg and a dose 35 times higher results practically in no tyramine potentiation, indicating the potential safety in clinical use.


TABLE 3

Potentiation of blood pressure response to tyramine	
Compound (5mg/kg)	Potentiation of blood pressure
(+)-6-Fluoropropargyl-1-aminoindan	1.2 fold
(+)-4-Fluoropropargyl-1-aminoindan	1.55 fold
(+)-5-Fluoropropargyl-1-aminoindan	2.25 fold

- Compounds were administered orally
- Tyramine was administered orally, 20mg/kg body weight

Claims

1. A compound of the formula




and pharmaceutically acceptable addition salts thereof.

2. A compound according to claim 1 selected from :

- 4-fluoro-N-propargyl-1-aminoindan or pharmaceutically acceptable acid addition salts thereof ;
- 5-fluoro-N-propargyl-1-aminoindan or pharmaceutically acceptable acid addition salts thereof ;
- 6-fluoro-N-propargyl-1-aminoindan or pharmaceutically acceptable acid addition salts thereof.

3. An optically pure (+) enantiomer of the compound of claim 1 or pharmaceutically acceptable addition salt thereof.


4. An optically pure (-) enantiomer of the compound of claim 1 or pharmaceutically acceptable addition salt thereof.
5. An optically pure (+) enantiomer of a compound according to claim 2.
6. A pharmaceutical composition comprising as active ingredient a compound as defined in claim 1 and a pharmaceutically acceptable carrier.
7. A pharmaceutical composition according to claim 6 in dosage unit form, each dosage unit containing from 1 to 20 mg of said active ingredient.
8. A pharmaceutical composition according to one of claims 6 or 7, additionally comprising levodopa and a decarboxylase inhibitor.
9. A pharmaceutical composition according to claims 7 and 8, wherein each dosage unit comprises 1-20 mg of the said active ingredient, 50-250 mg levopoda and 10-25 mg L-carbidopa.
10. A pharmaceutical composition according to claims 7 and 8, wherein each dosage unit comprises 2-10 mg of the said active ingredient, 50-250 mg levopoda and 12.5-50 mg benserazide.
11. Use of a compound according to any one of claims 1 to 5 for the manufacture of a pharmaceutical composition for the treatment of human patients suffering from Parkinson's disease, memory disorders, dementia of the Alzheimer type (DAT), depression or hyperactive syndrome in children.
12. A process for the preparation of a compound according to claim 1 comprising reacting racemic fluorinated 1-aminoindan of the formula



(II)

with propargyl bromide or propargyl chloride in the presence of an organic or inorganic base, optionally in the presence of a suitable solvent, and, if desired, converting the resulting free base of formula (I) in claim 1 into a pharmaceutically acceptable acid addition salt thereof.


13. A process for the preparation of a compound according to claim 3 comprising reacting the (-) enantiomer of fluorinated 1-aminoindan of the formula



(II)


with propargyl bromide or propargyl chloride in the presence of an organic or inorganic base, optionally in the presence of a suitable solvent, and if desired, converting the resulting (+) enantiomeric compound free base of formula (I) in claim 1 into a pharmaceutically acceptable acid addition salt thereof.

14. A process for the preparation of a compound according to claim 3 comprising reacting a racemic mixture of a compound of the formula



10 with propargyl bromide or propargyl chloride in the presence of an organic or inorganic base, optionally in the presence of a suitable solvent, and isolating the (+) enantiomeric form of the resulting racemic fluorinated 1-propargylaminoindan by chromatography, distillation, selective extraction, or conversion of the free base into a suitable acid addition salt and its fractional recrystallisation.


15 15. A method for the preparation of a (-) enantiomer of fluorinated 1-aminoindan of the formula



25 comprising selectively acylating a racemic mixture of said compound of formula (II) with a suitable acylating agent in the presence of the enzyme subtilisin A isolated from *Bacillus licheniformis* thereby forming a mixture comprising an acylated (+) enantiomer of fluorinated-1-aminoindan and a non-acylated (-) enantiomer of fluorinated-1-aminoindan, and separating the (-) fluorinated-1-aminoindan from the acylated (+) fluorinated-1-aminoindan by chromatography, distillation, selective extraction, or conversion of the free base into a suitable acid addition salt and its recrystallisation.

30 16. A method for the preparation of the (+)-fluoro-N-propargyl-1-aminoindan of claim 3 comprising:

35 (i) selectively acylating a racemic mixture of a compound of the formula



45 with a suitable acylating agent in the presence of the enzyme subtilisin A isolated from *Bacillus licheniformis* thereby forming a mixture comprising an acylated (+) enantiomer fluoro-1-aminoindan and a non-acylated (-) enantiomer of fluoro-1-aminoindan,

50 (ii) separating the (-) fluoro-1-aminoindan from the acylated (+) fluoro-1-aminoindan by chromatography, distillation, selective extraction, or conversion of the free base into a suitable acid addition salt and its fractional recrystallization, and

55 (iii) reacting the separated (-) fluoro-1-aminoindan with propargyl bromide or propargyl chloride in the presence of an organic or inorganic base, optionally in the presence of a suitable solvent, and isolating the (+) fluoro-N-propargyl-1-aminoindan formed as the free base form or in the form of a pharmaceutically acceptable acid addition salt thereof.


17. A fluorinated 1-aminoindan of formula (II) herein in racemate and enantiomeric forms and acid addition salts thereof.

18. A fluorinated 1-aminoindan according to claim 17 selected from:

- 4-fluoro-1-aminoindan in racemate and (-) enantiomer form and acid addition salts thereof;
- 5-fluoro-1-aminoindan in racemate and (-) enantiomer form and acid addition salts thereof;
- 6-fluoro-1-aminoindan in racemate, (-) enantiomer form and (+) enantiomer form and acid addition salts thereof.

Patentansprüche

1. Verbindung der Formel



und pharmazeutisch annehmbare Additionssalze derselben.

2. Verbindung nach Anspruch 1, ausgewählt aus:

- 4-Fluor-N-propargyl-1-aminoindan oder pharmazeutisch annehmbaren Säureadditionssalzen desselben;
- 5-Fluor-N-propargyl-1-aminoindan oder pharmazeutisch annehmbaren Säureadditionssalzen desselben;
- 6-Fluor-N-propargyl-1-aminoindan oder pharmazeutisch annehmbaren Säureadditionssalzen desselben.

3. Optisch reines (+)-Enantiomer der Verbindung nach Anspruch 1 oder pharmazeutisch annehmbares Additionssalz desselben.

4. Optisch reines (-)-Enantiomer der Verbindung nach Anspruch 1 oder pharmazeutisch annehmbares Additionssalz desselben.

5. Optisch reines (+)-Enantiomer einer Verbindung nach Anspruch 2.

6. Pharmazeutische Zusammensetzung, umfassend als wirksamen Bestandteil eine Verbindung wie in Anspruch 1 definiert und einen pharmazeutisch annehmbaren Träger.

7. Pharmazeutische Zusammensetzung nach Anspruch 6 in Form einer Dosiereinheit, wobei jede Dosiereinheit 1 bis 20 mg des wirksamen Bestandteils enthält.

8. Pharmazeutische Zusammensetzung nach irgendeinem der Ansprüche 6 oder 7, die zusätzlich Levodopa und einen Decarboxylase-Hemmer umfaßt.


9. Pharmazeutische Zusammensetzung nach den Ansprüchen 7 und 8, wobei jede Dosiereinheit 1-20 mg des wirksamen Bestandteils, 50-250 mg Levodopa und 10-25 mg L-Carbidopa umfaßt.

10. Pharmazeutische Zusammensetzung nach den Ansprüchen 7 und 8, wobei jede Dosiereinheit 2-10 mg des wirksamen Bestandteils, 50-250 mg Levodopa und 12,5-50 mg Benserazid umfaßt.

11. Verwendung einer Verbindung nach irgendeinem der Ansprüche 1 bis 5 zur Herstellung einer pharmazeutischen Zusammensetzung für die Behandlung von menschlichen Patienten, die an der Parkinson-Krankheit, Gedächtnisstörungen, Demenz vom Alzheimer-Typ (DAT), Depressionen oder, bei Kindern, an Hyperaktivitätssyndrom leiden.

12. Verfahren zur Herstellung einer Verbindung nach Anspruch 1, umfassend die Umsetzung eines racemischen fluo-


rierten 1-Aminoindans der Formel



(II)

mit Propargylbromid oder Propargylchlorid in Gegenwart einer organischen oder anorganischen Base, gegebenenfalls in Gegenwart eines geeigneten Lösungsmittels, und, falls gewünscht, die Überführung der resultierenden freien Base der Formel (I) in Anspruch in ein pharmazeutisch annehmbares Säureadditionssalz derselben.


13. Verfahren zur Herstellung einer Verbindung nach Anspruch 3, umfassend die Umsetzung des (-)-Enantiomers des fluorierten 1-Aminoindans der Formel



(II)

mit Propargylbromid oder Propargylchlorid in Gegenwart einer organischen oder anorganischen Base, gegebenenfalls in Gegenwart eines geeigneten Lösungsmittels, und, falls gewünscht, die Überführung der resultierenden freien Base der (+)-enantiomeren Verbindung der Formel (I) in Anspruch 1 in ein pharmazeutisch annehmbares Säureadditionssalz derselben.


14. Verfahren zur Herstellung einer Verbindung nach Anspruch 3, umfassend die Umsetzung einer racemischen Mischung einer Verbindung der Formel



(II)

mit Propargylbromid oder Propargylchlorid in Gegenwart einer organischen oder anorganischen Base, gegebenenfalls in Gegenwart eines geeigneten Lösungsmittels, und die Isolierung der (+)-enantiomeren Form des resultierenden racemischen fluorierten 1-Propargylaminoindans durch Chromatographie, Destillation, selektive Extraktion oder die Überführung der freien Base in ein geeignetes Säureadditionssalz und dessen fraktionierte Umkristallisation.

15. Verfahren zur Herstellung eines (-)-Enantiomers von fluoriertem 1-Aminoindan der Formel




(II)

umfassend die selektive Acylierung einer racemischen Mischung der Verbindung der Formel (II) mit einem geeigneten Acylierungsmittel in Gegenwart des aus *Bacillus licheniformis* isolierten Enzyms Subtilisin A, wodurch eine Mischung gebildet wird, umfassend ein acyliertes (+)-Enantiomer von fluoriertem 1-Aminoindan und ein nichtacyliertes (-)-Enantiomer von fluoriertem 1-Aminoindan, und die Trennung des (-)-Fluor-1-aminoindans vom acylierten (+)-Fluor-1-aminoindan durch Chromatographie, Destillation, selektive Extraktion oder Überführung der freien Base in ein geeignetes Säureadditionssalz und dessen Umkristallisation.

16. Verfahren zur Herstellung des (+)-Fluor-N-propargyl-1-aminoindans von Anspruch 3, umfassend:

(i) die selektive Acylierung einer racemischen Mischung einer Verbindung der Formel



(II)

mit einem geeigneten Acylierungsmittel in Gegenwart des aus *Bacillus licheniformis* isolierten Enzyms Subtilisin A, wodurch eine Mischung gebildet wird, umfassend ein acyliertes (+)-Enantiomer von Fluor-1-aminoindan und ein nichtacyliertes (-)-Enantiomer von Fluor-1-aminoindan,

(ii) die Trennung des (-)-Fluor-1-aminoindans vom acylierten (+)-Fluor-1-aminoindan durch Chromatographie, Destillation, selektive Extraktion oder die Überführung der freien Base in ein geeignetes Säureadditionssalz und dessen fraktionierte Umkristallisation, und

(iii) die Umsetzung des abgetrennten (-)-Fluor-1-aminoindans mit Propargylbromid oder Propargylchlorid in Gegenwart einer organischen oder anorganischen Base, gegebenenfalls in Gegenwart eines geeigneten Lösungsmittels, und die Isolierung des (+)-Fluor-N-propargyl-1-aminoindans, das in Form der freien Base oder in Form eines pharmazeutisch annehmbaren Säureadditionssalzes derselben gebildet wird.


17. Fluoriertes 1-Aminoindan der Formel (II) hierin in Racematform und den enantiomeren Formen sowie Säureadditionssalze desselben.

18. Fluoriertes 1-Aminoindan nach Anspruch 17, ausgewählt aus:

- 4-Fluor-1-aminoindan in der Racemat- und (-)-enantiomeren Form und Säureadditionssalzen desselben;
- 5-Fluor-1-aminoindan in der Racemat- und (-)-enantiomeren Form und Säureadditionssalzen desselben;
- 6-Fluor-1-aminoindan in der Racemat-, der (-)-enantiomeren Form und der (+)-enantiomeren Form und Säureadditionssalzen desselben.

Revendications

1. Composé de formule :



(I)

et ses sels d'addition pharmaceutiquement acceptables.

2. Composé selon la revendication 1, choisi parmi :

- le 4-fluoro-N-propargyl-1-aminoindane ou ses sels d'addition d'acide pharmaceutiquement acceptables,
- le 5-fluoro-N-propargyl-1-aminoindane ou ses sels d'addition d'acide pharmaceutiquement acceptables,
- le 6-fluoro-N-propargyl-1-aminoindane ou ses sels d'addition d'acide pharmaceutiquement acceptables.

3. Enantiomère (+) optiquement pur du composé de la revendication 1 ou d'un sel d'addition pharmaceutiquement acceptable de celui-ci.

4. Enantiomère (-) optiquement pur du composé de la revendication 1 ou d'un sel d'addition pharmaceutiquement acceptable de celui-ci.

5. Enantiomère (+) optiquement pur d'un composé selon la revendication 2.

6. Composition pharmaceutique comprenant comme ingrédient actif un composé tel que défini dans la revendication 1 et un support pharmaceutiquement acceptable.

7. Composition pharmaceutique selon la revendication 6 sous forme d'unités de dosage, chaque unité de dosage contenant de 1 à 20 mg dudit ingrédient actif.


8. Composition pharmaceutique selon l'une des revendications 6 et 7, comprenant en outre de la levodopa et un inhibiteur de la décarboxylase.

9. Composition pharmaceutique selon les revendications 7 et 8, dans laquelle chaque unité de dosage comprend 1 à 20 mg dudit ingrédient actif, 50 à 250 mg de levodopa et 10 à 25 mg de L-carbidopa.

10. Composition pharmaceutique selon les revendications 7 et 8, dans laquelle chaque unité de dosage comprend 2 à 10 mg dudit ingrédient actif, 50 à 250 mg de levodopa et 12,5 à 50 mg de benserazide.

11. Utilisation d'un composé selon l'une quelconque des revendications 1 à 5 pour la préparation d'une composition pharmaceutique pour le traitement de patients humains souffrant de la maladie de Parkinson, de troubles de la mémoire, de démence du type Alzheimer (DAT), de dépression ou de syndrome hyperactif chez les enfants.

12. Procédé de préparation d'un composé selon la revendication 1, qui comprend la réaction d'un 1-aminoindane fluoré racémique de formule




(II)

avec le bromure de propargyle ou le chlorure de propargyle en présence d'une base organique ou inorganique, éventuellement en présence d'un solvant approprié, et, si on le souhaite, la conversion de la base libre résultante

de formule (I) selon la revendication 1 en un sel d'addition d'acide pharmaceutiquement acceptable de celle-ci.


13. Procédé de préparation d'un composé selon la revendication 3, comprenant la réaction de l'énantiomère (-) d'un 1-aminoindane fluoré de formule



(II)

avec le bromure de propargyle ou le chlorure de propargyle en présence d'une base organique ou inorganique, éventuellement en présence d'un solvant approprié et, si on le souhaite, la conversion de la base libre du composé énantiomère (+) résultante de formule (I) selon la revendication 1 en un sel d'addition d'acide pharmaceutiquement acceptable de celle-ci.


14. Procédé de préparation d'un composé selon la revendication 3, comprenant la réaction d'un mélange racémique d'un composé de formule



(II)

avec le bromure de propargyle ou le chlorure de propargyle en présence d'une base organique ou inorganique, éventuellement en présence d'un solvant approprié, et l'isolement de la forme énantiomère (+) à partir du 1-propargylaminoindane fluoré racémique résultant par chromatographie, distillation, extraction sélective ou conversion de la base libre en un sel d'addition d'acide approprié et sa recrystallisation fractionnée.

15. Procédé de préparation d'un énantiomère (-) d'un 1-aminoindane fluoré de formule




(II)

comprenant l'acylation sélective d'un mélange racémique dudit composé de formule (II) avec un agent acylant approprié en présence de l'enzyme subtilisine A isolée à partir de *Bacillus licheniformis* pour former ainsi un mélange comprenant un énantiomère (+) acylé de 1-aminoindane fluoré et un énantiomère (-) non acylé de 1-aminoindane fluoré, et la séparation du (-) 1-aminoindane fluoré d'avec le (+) 1-aminoindane fluoré acylé par chromatographie, distillation, extraction sélective ou conversion de la base libre en un sel d'addition d'acide approprié et sa recrystallisation.

16. Procédé de préparation du (+)-fluoro-N-propargyl-1-aminoindane de la revendication 3 comprenant :

(i) l'acylation sélective d'un mélange racémique d'un composé de formule



(II)

avec un agent acylant approprié en présence de l'enzyme subtilisine A isolée à partir de *Bacillus licheniformis* pour former ainsi un mélange comprenant un énantiomère (+) acylé de fluoro-1-aminoindane et un énantiomère (-) non acylé de fluoro-1-aminoindane,

(ii) la séparation du (-)-fluoro-1-aminoindane d'avec le (+)-fluoro-1-aminoindane acylé par chromatographie, distillation, extraction sélective ou conversion de la base libre en un sel d'addition d'acide approprié et sa recristallisation fractionnée, et

(iii) la réaction du (-)-fluoro-1-aminoindane séparé avec le bromure de propargyle ou le chlorure de propargyle en présence d'une base organique ou inorganique, éventuellement en présence d'un solvant approprié, et l'isolement du (+)-fluoro-N-propargyl-1-aminoindane formé sous forme de la base libre ou sous forme d'un sel d'addition d'acide pharmaceutiquement acceptable de celle-ci.

17. 1-aminoindane fluoré de formule (II) sous forme de racémate et des énantiomères, et leurs sels d'addition d'acide.

18. 1-aminoindane fluoré selon la revendication 17 choisi parmi :

- le 4-fluoro-1-aminoindane sous forme de racémate et d'énantiomère (-) et leurs sels d'addition d'acide,
- le 5-fluoro-1-aminoindane sous forme de racémate et d'énantiomère (-) et leurs sels d'addition d'acide,
- le 6-fluoro-1-aminoindane sous forme de racémate, d'énantiomère (-) et d'énantiomères (+) et leurs sels d'addition d'acide.